

Density model for Clymene dolphin in the AFTT area - version 2

Laura Mannocci, Jason J Roberts, David L Miller, Patrick N Halpin

January 23, 2015

This report documents the density model developed for Clymene dolphin in the AFTT area. It provides information on available data, methodological decisions, the selected model, predictions, uncertainty, model checking and qualitative evaluation of predictions based on the literature. Information on classification of ambiguous sightings, detection function fitting and $g(0)$ estimates can be found in the EEZ model report for this taxon (Roberts et al. 2015).

Citation for this model: Mannocci L, Roberts JJ, Miller DL, Halpin PN (2015) Density model for Clymene dolphin in the AFTT area. Version 2, 2015-01-23. Marine Geospatial Ecology Lab, Duke University, Durham, NC.

Citation for the related peer-review publication: Mannocci L, Roberts JJ, Miller DL, Halpin PN. Here be dragons: extrapolating cetacean densities into the unsurveyed high seas of the western North Atlantic. Submitted to Ecological Applications.

1- Available data

Table 1: Effort (km) and sightings per region (CAR: Caribbean, EC: East coast, EU: European Atlantic, GM: Gulf of Mexico, MAR: Mid-Atlantic ridge).

Region	Effort	Sightings
EC	1044357.7	11
GOM	194715.3	78
All regions	1239073.1	89

Table 2: Effort (km) and sightings per month.

Month	Effort	Sightings
January	77450.08	1
February	113657.63	4
March	111498.11	2
April	117481.09	10
May	149765.03	37
June	129393.69	9
July	135693.85	15
August	129660.43	7
September	70592.28	3
October	76650.01	1
November	69210.92	0
December	58019.93	0
All Months	1239073.05	89




Figure 1: Map of segments (black lines) and sighting locations (red dots). An Albers equal area projection optimized for the AFT area is used.

2- Methodological decisions

Modeled taxon

Clymene dolphin (*Stenella clymene*)

Model type

Due to the small sample size, we fitted a simple habitat-based density model with sea surface temperature (SST) as the single covariate for this taxon. SST was successfully used in predictive distribution models of cetaceans in data poor situations (Kaschner et al. 2006).

Modeled season

We fitted a year-round model as we found no evidence in the literature that this taxon undertakes extensive migrations or exhibits contrasting behaviors (e.g. feeding versus breeding) in different seasons.

Segments

We used segments from the east coast, Gulf of Mexico, and Caribbean. Although segments from the Caribbean did not include sightings, their incorporation prevented an increase of the extent of environmental extrapolation and resulted in a negligible drop in the percentage of presence segments (from 0.072% to 0.071% after adding the Caribbean segments).

Temporal resolution of predictions

Since there was insufficient evidence in the literature to support the monthly variations in predicted densities, we produced a year-round density prediction by averaging the twelve monthly density predictions.

3- Best model

- Selected covariates: sea surface temperature
- Model summary:

```
##
## Family: Tweedie(p=1.558)
## Link function: log
##
## Formula:
## abundance ~ s(SST, k = 4, bs = "ts") + offset(log(area_km2))
## <environment: 0x067e03bc>
##
## Parametric coefficients:
##           Estimate Std. Error t value Pr(>|t|)
## (Intercept) -10.774      1.544  -6.978 3.01e-12 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Approximate significance of smooth terms:
##           edf Ref.df    F  p-value
## s(SST) 2.763     3 11.26 7.31e-08 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## R-sq.(adj) = 0.000306  Deviance explained = 43.6%
## -REML = 1347  Scale est. = 1555.2    n = 124752
```






Figure 2: GAM term plots with the log-transformed abundance on the y axis. The solid green line is the smooth function fitted to the data. The solid red line is the smooth function extrapolated to all covariate values in the prediction area. The dashed lines represent the approximate 95% confidence intervals. The rug plot on the x-axis shows the range of covariate values sampled in the data. Note that transformations were used for some covariates.

4- Environmental envelopes


January




February




March




April



May



June






Figure 3: Monthly environmental envelopes for CMC_SST. White cells within the AFTT polygon indicate areas where covariate values fell beyond the range of covariate values sampled by the surveys.

5- Predictions




Figure 4: Mean predicted densities (individuals 100 km⁻²) in the AFTT area. Areas where we extrapolated beyond the sampled covariate ranges are indicated with black crosshatches. An Albers equal area projection is used.

Table 3: Mean predicted abundance (individuals) in the AFTT area and associated coefficient of variation (CV). The CV only reflects uncertainty in the estimated GAM parameters. It does not consider extrapolation beyond the sampled covariate ranges and is therefore strongly underestimated.

Abundance	CV
203211	0.173

6- Uncertainty




Figure 5: Mean predicted coefficients of variation (unit-less) in the AFTT area. Areas where we extrapolated beyond the sampled covariate ranges are indicated with black crosshatches. An Albers equal area projection is used.

7- Residual diagnostics




Figure 6: Diagnostic plots of deviance residuals. The normal Q Q plot is useful to assess goodness of fit.




Figure 7: Diagnostic plots of randomized quantile residuals. Randomized quantile residuals (exactly normal residuals) are the most adapted residuals to visualize diagnostic plots of regression models applied to count data. The plots of residuals versus linear predictor and response versus fitted values are useful to investigate patterns in the residuals (e.g. non constant variance).




Figure 8: Map showing the spatial distribution of deviance residuals with positive residuals in red and negative residuals in blue.

8- Qualitative evaluation of predictions

Model predictions are consistent with the reported distribution of Clymene dolphins throughout tropical and warm temperate waters of the Atlantic Ocean, Caribbean Sea, and Gulf of Mexico (Fertl et al. 2003). Predictions in the southern Gulf of Mexico are compatible with sightings collected from opportunistic shipboard surveys (Ortega-Ortiz 2002).

Future model improvements

The acquisition of additional sightings data would increase the sample size and potentially allow fitting a habitat-based density model with a full variable selection procedure.

REFERENCES

Fertl, D., T. A. Jefferson, I. B. Moreno, A. N. Zerbini, and K. D. Mullin. 2003. Distribution of the Clymene dolphin *Stenella clymene*. *Mammal Review* 33:253-271.

Kaschner, K., R. Watson, A. W. Trites, and D. Pauly. 2006. Mapping world-wide distributions of marine mammal species using a relative environmental suitability (RES) model. *Marine Ecology Progress Series* 316:285-310.

Ortega-Ortiz, J. 2002. Multiscale analysis of cetacean distribution in the Gulf of Mexico. PhD dissertation, Texas A&M University.